



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by H. C. WHITAKER, C. E., Ph. D., Manual Training School, Philadelphia, Pa.; MARCUS BAKER, U. S. Geological Survey, Washington, D. C.; and G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.

Plotting the curves represented by the two equations, they are seen to intersect at (0, 1) and also in the neighborhood of (.6, .8).

$$\frac{\log(7-6^y)}{\log 5} - \frac{\log(4-3^y)}{\log 2} = 0.$$

By Double Position the roots near .6, .8 are found to be $x=.565585$, $y=.841307$; these with the values $x=0$, $y=1$ seem to be the only real roots.

Also solved by JOHN G. KELLER, State Normal School, Albany, N. Y.
The Proposer sent results only, to-wit: $x=.56558312$, $y=.8413092$.

GEOMETRY.

123, Proposed by P. C. CULLEN, Indianola, Iowa.

If the bisectors of the base angles of a triangle are equal the triangle is isosceles.

I. Direct Demonstration by G. I. HOPKINS, A. M., Professor of Physics and Astronomy, High School, Manchester, N. H.

CONSTRUCTION. Make $\angle BDK = \angle FAB$, and draw through B the line NK , making $\angle DBK = \angle AFB$. Draw the perpendiculars KP and AN , and join A and K .

DEMONSTRATION. Since angles FAB and AFB are two angles of a triangle, DK and NK will meet.

\therefore triangles DBK and AFB are equal.

$\therefore DK = AB$ and $BK = BF$.

$\angle AHB = \angle ADH + \angle DAH = \angle ADH + \angle BDK$.

$\therefore \angle AHB = \angle ADK$. Also $\angle AHB = \angle HBF + \angle HFB = \angle HBA + \angle HBK$.

$\therefore \angle AHB = \angle ABK$. $\therefore \angle ADK = \angle ABK$.

$\therefore \angle KDP = \angle ABN$. $\therefore \triangle DPK = \triangle ABN$.

$\therefore AN = KP$, and $NB = DP$.

$\therefore \triangle APK = \triangle AKN$.

$\therefore AP = NK$, and $\therefore AD = BK$. $\therefore AB = FB$. $\therefore \triangle ADK = \triangle AFB$.

$\therefore \angle ABD = \angle FAB$. $\therefore \angle DAB = \angle AFB$.


$\therefore AC = BC$, and the triangle ABC is isosceles.

Q. E. D.

II. Demonstration by L. E. DICKSON, Ph. D., Assistant Professor of Mathematics, University of Chicago, Chicago, Ill.

Let d_a be the length of the bisector of the angle A opposite to side a . Expressing the area of the given triangle and the areas of the triangles into which it is divided by the bisector of angle A , we get

$$\frac{1}{2}bc\sin A = \frac{1}{2}bd_a\sin\frac{1}{2}A + \frac{1}{2}cd_a\sin\frac{1}{2}A.$$



$$\therefore d_a = \frac{2bc}{b+c} \cos \frac{1}{2} A = \frac{2\sqrt{[s(s-a)bc]}}{b+c}, \text{ where } 2s = a + b + c.$$

$$\text{Hence, if } d_a = d_b, \frac{2\sqrt{[s(s-a)bc]}}{b+c} = \frac{2\sqrt{[s(s-b)ac]}}{a+c}.$$

$$\therefore 2(a+c)^2(s-a)b = 2(b+c)^2(s-b)a.$$

$$\therefore (a+c)^2(b+c-a)b - (b+c)^2(a+c-b)a = 0.$$

$$\therefore (a-b)[c^3 + c^2(a+b) + 3abc + ab(a+b)] = 0.$$

Since the second factor is positive, it does not vanish. Hence $a = b$.

We have finally received an elementary direct proof of this theorem. The proof by Professor Hopkins is, I believe, without a flaw, and is the proof so long sought for by a number of mathematicians, among whom was Isaac Todhunter. This demonstration of Professor Hopkins' was examined by one of the ablest mathematicians in this country, and was pronounced by him to be correct. The trigonometric proof by Dr. Dickson is also flawless. We are glad to publish both of these proofs since the demonstration which we gave in Vol. VII, page 223, has been assailed.

Last January a year ago, we received an indirect proof from Prof. A. Anderson, of the University of North Carolina. Professor Anderson's demonstration is free from error and is substantially the proof given in Dr. Halsted's Synthetic Geometry, page 44, though Professor Anderson's figure is drawn quite differently from the one in Dr. Halsted's Geometry.

We shall be pleased to receive, from our readers, opinions on the above demonstrations.

153. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, O.

If P, P', Q, Q' be the extremities of two chords of a conic section, and both chords pass through the point A , show that the sum of the squares of the reciprocals of AP, AP', AQ, AQ' is constant.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

$$\frac{1}{AP} = \frac{1 + e \cos \theta}{l}, \quad \frac{1}{AP'} = \frac{1 + e \cos(\pi + \theta)}{l} = \frac{1 - e \cos \theta}{l}.$$

$$\frac{1}{AQ} = \frac{1 + e \cos(\frac{1}{2}\pi + \theta)}{l} = \frac{1 - e \sin \theta}{l}.$$

$$\frac{1}{AQ'} = \frac{1 + e \cos(3\pi/2 + \theta)}{l} = \frac{1 + e \sin \theta}{l}.$$

$$\begin{aligned} & \therefore \frac{1}{(AP)^2} + \frac{1}{(AP')^2} + \frac{1}{(AQ)^2} + \frac{1}{(AQ')^2} \\ &= \frac{(1 + e \cos \theta)^2 + (1 - e \cos \theta)^2 + (1 - e \sin \theta)^2 + (1 + e \sin \theta)^2}{l^2} = \frac{2(2 + e^2)}{l^2} = \text{a constant.} \end{aligned}$$

158. Proposed by JOHN MACNIE, A. M., Professor of Latin, University of North Dakota.

Show by a simple diagram that:

(a) If the angle-sum of an equilateral triangle is constant, that constant is a straight angle.